

Getting Started With the Orenitram Titration Kit



A guide to helping you confidently start your treatment with Orenitram® (treprostinil) Extended-Release Tablets.

Please see Important Safety Information throughout and on back, and the Full Prescribing Information and Patient Information for Orenitram in pocket.


orenitram®
treprostinil
EXTENDED-RELEASE TABLETS

Why Orenitram?


Orenitram works differently than other oral medications you may already be taking for your PAH. Orenitram mimics the effects of prostacyclin, a natural substance found in your body that opens up blood vessels so it's easier for your heart to pump blood through your lungs.

Together, you and your healthcare provider have made the important decision to add Orenitram to your current treatment plan because it has been proven to help fight your PAH—helping you achieve your treatment goals.

Reaching your target dose to achieve treatment goals

Reaching your target dose is important because it may help you achieve the most out of treatment with Orenitram. You'll get there through **titration**, the process of increasing your dosage until you reach the dose that is appropriate for you.

The Orenitram Titration Kit follows a 3-month titration plan, putting you on the path toward reaching your target dose.



PAH=pulmonary arterial hypertension.

IMPORTANT SAFETY INFORMATION for Orenitram

Who should not take Orenitram?

Do not take Orenitram if you have severe liver problems.

What should I tell my healthcare provider before taking Orenitram?

Tell your healthcare provider:

- If you have liver problems or diverticulosis.

Please see Important Safety Information throughout and on back, and the Full Prescribing Information and Patient Information for Orenitram in pocket.

The Orenitram Titration Kit— designed for a simple, confident start

We know that starting a new treatment isn't easy—that's why the all-in-one Orenitram Titration Kit was created to make your first 3 months on Orenitram as straightforward as possible.



3 MONTHLY KITS

Each Monthly Kit includes:

- Support based on where you are during titration
- A simple outline of that month's Orenitram doses

4 WEEKLY BOXES PER MONTH

Each Weekly Box includes:

- How to take Orenitram
- What to do if you miss a dose

7 DAILY PACKS PER WEEK

Each Daily Pack includes:

- Your exact morning, afternoon, and evening doses of Orenitram
- Tools to help ensure you're staying on track



Watch this video to see how the Orenitram Titration Kit works.



IMPORTANT SAFETY INFORMATION for Orenitram

What should I tell my healthcare provider before taking Orenitram? (continued)

Tell your healthcare provider:

- If you are pregnant, breastfeeding, and/or plan to become pregnant or breastfeed. It is not known if Orenitram will harm your unborn baby or if Orenitram passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with Orenitram.
- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Orenitram and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your healthcare provider. Especially tell your healthcare provider if you take another medicine that contains treprostinil, such as Remodulin® or Tyvaso®.



Taking Orenitram

Talk with your healthcare provider about food choices and combinations that may work best for you.

You'll take Orenitram with food 3 times a day, once every 8 hours.*



Expected adverse events and how to manage them

Because of the way Orenitram works, you may experience side effects before you see the benefits of therapy. **Before you start therapy, create a plan with your doctor to manage these expected effects so you can commit to treatment and find the dose your body needs to feel better and fight your PAH.**

Below are some of the most common adverse events associated with Orenitram—and suggestions for managing them:

Headaches

- ✓ Acetaminophen
- ✓ Ibuprofen

Nausea and vomiting

- ✓ Taking with food
- ✓ Anti-nausea and anti-vomiting medicine

Diarrhea

- ✓ Adding fiber to your diet
- ✓ Anti-diarrhea medicine

United Therapeutics does not provide medical advice. Your healthcare provider will work with you to develop a plan to manage side effects.



Learn more about side effects and how to manage them by visiting www.orenitram.com/side-effects.

*Patients who are not on the Titration Kit may also take Orenitram 2 times a day, or once every 12 hours.

IMPORTANT SAFETY INFORMATION for Orenitram

How should I take Orenitram?

- **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**
- Orenitram is usually taken 3 times a day (about every 8 hours) or 2 times a day (about every 12 hours). Your healthcare provider will tell you how often you should take Orenitram. If you have side effects, your healthcare provider may tell you to change your dose or when you take Orenitram. Take Orenitram with food.

Tips for a successful start with Orenitram



Work with your doctor to learn about expected side effects and commit to a plan to proactively help manage them.



Identify, discuss, and revisit your personal goals to help you fight through hard days—goals such as taking walks, doing yardwork, or running errands.



Stay strong. Remember your goals and how important it is to stay committed to achieving your target dose.

- After the first 3 months on the Titration Kit, you will receive Orenitram tablets in pill bottles and can continue your routine



To anybody starting therapy with Orenitram, give yourself time, remember your goals, listen to your body, and go with what you can do.

— an Orenitram patient

IMPORTANT SAFETY INFORMATION for Orenitram

How should I take Orenitram? (continued)

- Swallow Orenitram tablets whole. Do not split, chew, crush, or break your Orenitram tablets. Do not take Orenitram tablets that are damaged or broken. **If Orenitram tablets are not taken whole, they may release too much medicine at one time. This can lead to side effects.**
- If you miss your dose of Orenitram, take the missed dose as soon as possible with food.
- If you miss 2 or more doses of Orenitram, call your healthcare provider to see if you need to change your dose.

Please see Important Safety Information throughout and on back, and the Full Prescribing Information and Patient Information for Orenitram in pocket.



Support for your treatment journey with Orenitram



1-877-242-2738



Accredo Health Group, Inc.
an Express Scripts company

1-866-344-4874



Specialty Pharmacy (SP) Support

Specialty Pharmacies help you with medication and supplies, in-home training, follow-up nurse visits, and have a 24-hour hotline in case anything comes up. United Therapeutics works with 2 SPs to support Orenitram patients.

United Therapeutics ASSIST

ASSIST helps eligible patients with therapy access, insurance coverage, and financial assistance information.



Co-Pay Assistance Program

With our co-pay program, eligible patients may pay as little as \$0 per prescription for Orenitram.



Virtual Patient Educator

United Therapeutics' Archways program is a virtual, educational program that offers helpful disease information and treatment resources. **Archways is available at no cost to you and your caregivers.** To begin, follow one of these three simple steps:

- Call 1-888-327-2492
- Text "START" to 882724 (Standard data rates may apply)
- Visit www.UTArchways.com



Learn more and hear from other patients about their Orenitram experience by visiting www.orenitram.com.

IMPORTANT SAFETY INFORMATION for Orenitram

How should I take Orenitram? (continued)


- If you take too much Orenitram, call your healthcare provider or go to the nearest hospital emergency room right away.
- You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.

Frequently asked questions about titrating with Orenitram



Q. What is titration with Orenitram?

A. Titration is the process of slowly increasing your dose of Orenitram over time until you find the most effective dose for you.



Q. How does the Orenitram Titration Kit help me start my treatment?

A. Starting a new treatment can be challenging, but the Titration Kit is designed to make treatment with Orenitram simpler. Instead of receiving your medication in bottles, your first 3 months of Orenitram will be organized each day in packs.

The Titration Kit contains helpful information about dosing, simple instructions, and reminders to know your exact morning, afternoon, and evening doses of Orenitram.



Q. What happens if I miss a dose of Orenitram?

A. If you've started on Orenitram and you miss a dose, take the dose as soon as possible with food. If you miss 2 or more doses, call your healthcare provider for guidance.

Do not stop taking Orenitram without consulting your healthcare provider.



If you have any other questions, see the Orenitram Titration Kit or visit www.orenitram.com/kit.



IMPORTANT SAFETY INFORMATION for Orenitram

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- Stopping Orenitram suddenly may cause worsening of your PAH symptoms. **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**

Please see Important Safety Information throughout and on back, and the Full Prescribing Information and Patient Information for Orenitram in pocket.



IMPORTANT SAFETY INFORMATION for Orenitram

What is Orenitram?

Orenitram is a prescription medicine used to treat pulmonary arterial hypertension (PAH) which is high blood pressure in the arteries of your lungs. Orenitram can help slow down the progression of your disease and improve your ability to exercise. It is not known if Orenitram is safe and effective in children.

Who should not take Orenitram?

Do not take Orenitram if you have severe liver problems.

What should I tell my healthcare provider before taking Orenitram?

Tell your healthcare provider:

- If you have liver problems or diverticulosis.
- If you are pregnant, breastfeeding, and/or plan to become pregnant or breastfeed. It is not known if Orenitram will harm your unborn baby or if Orenitram passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with Orenitram.
- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Orenitram and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your healthcare provider. Especially tell your healthcare provider if you take another medicine that contains treprostinil, such as Remodulin® or Tyvaso®.

How should I take Orenitram?

- **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**
- Orenitram is usually taken 3 times a day (about every 8 hours) or 2 times a day (about every 12 hours). Your healthcare provider will tell you how often you should take Orenitram. If you have side effects, your healthcare provider may tell you to change your dose or when you take Orenitram. Take Orenitram with food.
- Swallow Orenitram tablets whole. Do not split, chew, crush, or break your Orenitram tablets. Do not take Orenitram tablets that are damaged or broken. **If Orenitram tablets are not taken whole, they may release too much medicine at one time. This can lead to side effects.**
- If you miss your dose of Orenitram, take the missed dose as soon as possible with food.
- If you miss 2 or more doses of Orenitram, call your healthcare provider to see if you need to change your dose.
- If you take too much Orenitram, call your healthcare provider or go to the nearest hospital emergency room right away.
- You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- Stopping Orenitram suddenly may cause worsening of your PAH symptoms. **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**
- **The most common side effects of Orenitram include** headache, diarrhea, nausea, vomiting, flushing, and pain in arms, legs, and jaw. These are not all of the possible side effects of Orenitram. Tell your healthcare provider if you have any side effect that bothers you or does not go away.
- Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at www.fda.gov/MedWatch or call 1-800-FDA-1088.

The risk information provided here is not comprehensive. To learn more about Orenitram, talk with your healthcare provider. Please see Full Prescribing Information and Patient Information at www.orenitram.com or call Customer Service at 1-877-UNITHER (1-877-864-8437).

OREISIconOCT19



Orenitram is a registered trademark of United Therapeutics Corporation.

All other brands are trademarks of their respective owners. The makers of these brands are not affiliated with and do not endorse United Therapeutics or its products.

© 2023 United Therapeutics Corporation. All rights reserved. US/ORE/0644 Printed in USA.

