

SOOD CAPTURE MORE SOOD CAPTURE MORE SOOD CAPTURE MORE SOOT CAPTURE

Adding Orenitram is proven to delay the progression of PAH and may help improve symptoms—to help you do more and help control your PAH

IMPORTANT SAFETY INFORMATION for Orenitram

Who should not take Orenitram?

Do not take Orenitram if you have severe liver problems.

Please see complete Important Safety Information throughout this brochure and Full Prescribing Information and Patient Information for Orenitram at www.orenitram.com. PAH=pulmonary arterial hypertension.

orenitram° treprostinil



PAH can make you miss out on the good days you want to capture...

Is it time to consider adding Orenitram?

PAH symptoms such as shortness of breath or feeling tired may mean you just can't do as many activities in your day-to-day life. Because PAH is a progressive disease, you may find yourself becoming less active over time.

With Orenitram, you may be able to do more of the things that matter to you

Your current medicine for PAH may be helping your symptoms, but you should know that there may be more you can be doing.

Orenitram has been proven to delay the progression of PAH and may help improve symptoms.

Now is the time to work with your healthcare provider to determine if Orenitram is right for you.

What is Orenitram?

Orenitram is a prescription medicine used to treat pulmonary arterial hypertension (PAH) which is high blood pressure in the arteries of your lungs. Orenitram can help slow down the progression of your disease and improve your ability to exercise. It is not known if Orenitram is safe and effective in children.



GG My 6-minute walk distance was not improving. My doctor recommended starting Orenitram to help me do more physically and feel better in my daily life.

- Angelie, an Orenitram patient

IMPORTANT SAFETY INFORMATION for Orenitram What should I tell my healthcare provider before taking Orenitram? Tell your healthcare provider:

• If you have liver problems or diverticulosis.



How does Orenitram work?

Orenitram is an oral medicine that works differently from other oral medicines you may already be taking for PAH. That's why adding it to your treatment plan may be an important next step to help you do more and help control your PAH.



PAH affects 3 substances in the body

PAH symptoms can occur when the blood vessels inside your lungs become narrow, making it harder for blood to flow. These changes are thought to be caused by imbalances of natural substances in 3 pathways in your body. There are medicines that treat each of those 3 pathways.

NITRIC OXIDE

o
ENDOTHELIN
o
PROSTACYCLIN


Depending on which pathway you are treating, there may be more you can do. Adding Orenitram may help make sure you don't miss out on treating the prostacyclin pathway.

IMPORTANT SAFETY INFORMATION for Orenitram What should I tell my healthcare provider before taking Orenitram? (cont) Tell your healthcare provider:

• If you are pregnant, breastfeeding, and/or plan to become pregnant or breastfeed. It is not known if Orenitram will harm your unborn baby or if Orenitram passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with Orenitram.

Why is the prostacyclin pathway important?

When you have PAH, your body may not produce enough prostacyclin, a natural substance that keeps your blood vessels open and working properly.



the effects of natural prostacyclin, helping to open up blood vessels so it's easier for your heart to pump blood through your lungs.

Orenitram helps to replace the natural prostacyclin you may be missing.

Talk to your healthcare provider to see if Orenitram may help you reach your treatment goals so you can start to capture more good days.

IMPORTANT SAFETY INFORMATION for Orenitram What should I tell my healthcare provider before taking Orenitram? (cont) Tell your healthcare provider:

 About all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Orenitram and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your healthcare provider. Especially tell your healthcare provider if you take another medicine that contains treprostinil, such as Remodulin® or Tyvaso®.



Orenitram—helping you capture more good days

Orenitram is the tablet form of the same proven and trusted type of medicine that has been available as a pump therapy for more than 18 years. Adding Orenitram may help you do more and help control your PAH.

Proven to delay disease progression

In a clinical study, patients who added Orenitram were...

61% LESS LIKELY

TO HAVE THEIR PAH PROGRESS OVER TIME COMPARED WITH THOSE NOT TAKING ORENITRAM*

*In a clinical study that included 690 patients who were all taking a single oral PAH medicine (PDE-5i, sGCS, or ERA). At the beginning of the study, these patients added either Orenitram or placebo to their treatment.

IMPORTANT SAFETY INFORMATION for Orenitram

How should I take Orenitram?

 Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.




ERA=endothelin receptor antagonist; PDE-5i=phosphodiesterase type 5 inhibitor; sGCS=soluble guanylate cyclase stimulator.

[†] In a clinical study that included 349 patients not taking any PAH therapy. At the beginning of the study, these patients began taking either Orenitram or placebo.

Orenitram may help you do more and help control your PAH

In clinical studies, adding Orenitram helped patients improve in several ways:



IMPORTANT SAFETY INFORMATION for Orenitram

How should I take Orenitram? (cont)

Orenitram is usually taken 3 times a day (about every 8 hours) or 2 times a
day (about every 12 hours). Your healthcare provider will tell you how often
you should take Orenitram. If you have side effects, your healthcare provider
may tell you to change your dose or when you
take Orenitram. Take Orenitram with food.

Please see complete Important Safety Information throughout this brochure and Full Prescribing Information and Patient Information for Orenitram at www.orenitram.com.

FXTENDED-RELEASE TABLETS

Orenitram may help improve your risk status*





Patients taking Orenitram either improved or maintained their risk status—and improvement was seen as early as 3 months. In fact, more than 30% of patients improved and were able to maintain improvement through week 60.

Why is risk status important?

Your healthcare provider may use risk status to make decisions about your treatment. It helps them determine how you're doing today and predict how you will be doing in the future.

Start the conversation

Find a helpful discussion guide at the end of this brochure, with questions you can ask your healthcare provider about adding Orenitram to your treatment. Fill it out and bring it with you to your next visit.



*In a clinical study that included 690 patients who were all taking a single oral PAH medicine (PDE-5i, sGCS, or ERA). At the beginning of the study, these patients added either Orenitram or placebo to their treatment.

IMPORTANT SAFETY INFORMATION for Orenitram How should I take Orenitram? (cont)

- Swallow Orenitram tablets whole. Do not split, chew, crush, or break your Orenitram tablets. Do not take Orenitram tablets that are damaged or broken. If Orenitram tablets are not taken whole, they may release too much medicine at one time. This can lead to side effects.
- If you miss your dose of Orenitram, take the missed dose as soon as possible with food.

GG With Orenitram, I'm able to plan more and make more good memories. I can share even the simple things, like cooking or laughing, with my family and friends.

- Angelie, an Orenitram patient

IMPORTANT SAFETY INFORMATION for Orenitram How should I take Orenitram? (cont)

- If you miss 2 or more doses of Orenitram, call your healthcare provider to see if you need to change your dose.
- If you take too much Orenitram, call your healthcare provider or go to the nearest hospital emergency room right away.



Getting started with Orenitram

Like other medicines you may be taking, Orenitram is an oral tablet. Orenitram is taken with food, usually 3 times daily (every 8 hours).*



Your healthcare provider will tell you exactly how to take Orenitram, and together, you can create a plan that fits into your daily life.

Personalized dosing

Since PAH is a progressive disease, your treatment needs may change over time. With Orenitram, you may not have to start a new medication. Your healthcare provider can continue to increase or change your dose to meet your needs so you may be able to stay with Orenitram for the long term to help control your PAH.[†]

Orenitram is available in 5 tablet strengths, making it easier for your healthcare provider to adjust your dose.



Available in 5 tablet strengthsNot shown at actual size.

IMPORTANT SAFETY INFORMATION for Orenitram

How should I take Orenitram? (cont)

• You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.

^{*}Orenitram may also be taken 2 times a day (every 12 hours). Tablet strength may be increased depending on how well you tolerate treatment.

[†]There is no maximum dose for Orenitram, so your healthcare provider can continue to increase your dose as long as you are able to tolerate the medication.



Knowing what to expect from treatment

Orenitram may cause side effects. The most common side effects with Orenitram include headache, diarrhea, nausea, vomiting, flushing, and pain in the arms, legs, and jaw.

Your healthcare provider will talk to you about potential side effects so you can feel prepared and plan ahead to help address them.

IMPORTANT SAFETY INFORMATION for Orenitram What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

 Stopping Orenitram suddenly may cause worsening of your PAH symptoms. Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.



Support from the start



If you and your healthcare provider decide that Orenitram may help you reach your treatment goals, United Therapeutics, the maker of Orenitram, provides support at every step in the process.



Assistance from a Specialty Pharmacy

When you take Orenitram, you don't get it from your local pharmacy. United Therapeutics works with Specialty Pharmacies (SP) that specialize in PAH to supply your medication.

Your SP will help you at each step of your treatment with Orenitram, such as:

- Scheduling nurse visits to get you started with treatment
- Offering ongoing assistance through in-person or virtual visits and 24-hour phone support
- Coordinating the delivery of your medicine



ASSIST is a team at United Therapeutics committed to helping you with your prescription, from answering questions about your prescription to explaining your insurance coverage, and much more. To learn more, visit **www.orenitram.com**.

IMPORTANT SAFETY INFORMATION for Orenitram

What are the possible side effects of Orenitram? (cont)

• The most common side effects of Orenitram include headache, diarrhea, nausea, vomiting, flushing, and pain in arms, legs, and jaw. These are not all of the possible side effects of Orenitram. Tell your healthcare provider if you have any side effect that bothers you or does not go away.

DISCUSSION GUIDE

Talking to your healthcare provider about Orenitram



Cost savings

With our co-pay program, eligible patients may pay as little as \$5 per prescription for Orenitram.* Learn more at **www.UTcopay.com**.



Orenitram Connect

To learn more about adding Orenitram to your PAH treatment plan, join Orenitram Connect. With this program, you'll receive e-newsletters, helpful information, and other useful tools. Sign up at www.orenitram.com.

*TO ENROLL IN THIS PROGRAM, YOU MUST UNDERSTAND AND AGREE TO COMPLY WITH THE ELIGIBILITY REQUIREMENTS AND TERMS OF USE.

Eligibility requirements for this program are:

Patients must be 18 years or older to use this Program. The Program is valid only for patients with commercial (also known as private) insurance who are taking the medication for an FDA-approved indication. Patients using Medicare, Medicaid, or any other state or federal government program to pay for their medications are not eligible. Patients who start utilizing government coverage during the term of the Program will no longer be eligible. Eligible patients must be residents of the United States or Puerto Rico.

For full program details and Terms and Conditions, visit www.UTcopay.com.

IMPORTANT SAFETY INFORMATION for Orenitram What are the possible side effects of Orenitram? (cont)

 Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at www.fda.gov/MedWatch or call 1-800-FDA-1088.





Capture more good days with Orenitram

Orenitram may be just what you need to reach your treatment goals

- Orenitram is proven to delay the progression of PAH and may help reduce symptoms that can impact your day-to-day life
- Personalized dosing means you may stay with
 Orenitram for the long term to help control your PAH*
- United Therapeutics offers support throughout your treatment journey with Orenitram

*There is no maximum dose for Orenitram, so your healthcare provider can continue to increase your dose as long as you are able to tolerate the medication.

Ask your healthcare provider if adding Orenitram may help you do more and help control your PAH.



To learn more about Orenitram, visit www.orenitram.com.

IMPORTANT SAFETY INFORMATION for Orenitram

The risk information provided here is not comprehensive. To learn more about Orenitram, talk with your healthcare provider. Please see Full Prescribing Information and Patient Information at www.orenitram.com or call Customer Service at 1-877-UNITHER (1-877-864-8437).

OREISIconOCT19

Please see complete Important Safety Information throughout this brochure and Full Prescribing Information and Patient Information for Orenitram at www.orenitram.com.

Orenitram is a registered trademark of United Therapeutics Corporation. © 2020 United Therapeutics Corporation. All rights reserved. US/ORE/0465



DISCUSSION GUIDE

Talking to your healthcare provider about Orenitram

Your healthcare team is your best source of information about your treatment options. Having open, honest discussions with them about your treatment goals can help determine if Orenitram may be right for you. Included below are some things to think about and questions you can ask about adding Orenitram to your treatment plan.

First, think about whether you are getting the most out of your current treatment and if it might be time to do more to help you feel better.

Fill in or write down what your goals are for treatment and your future (for example, taking a family vacation or being able to do more daily errands):



Next, ask your healthcare provider the following questions and fill in or write down their responses:

| questions and fill in or write down their responses. |
|--|
| How may Orenitram help me meet my treatment goals? |
| How may Orenitram help improve my symptoms? |
| How is Orenitram different from my current medicine? |
| What can I expect from treatment? |
| Talk to your healthcare provider to see if Orenitram is right for you. |